

LHe/WYo

22 March 2024

Dear parent/carer

Increase in Scarlet Fever

We are writing to provide some information about the increase in scarlet fever which is above expected levels for the time of year.

We would like to take this opportunity to remind you of the signs, symptoms and the actions to be taken if you think that you or your child might have scarlet fever.

Signs and symptoms of scarlet fever

Scarlet fever is a common childhood infection caused by Group A streptococci. It is not usually serious, but should be treated with antibiotics to reduce the risk of complications (such as pneumonia) and spread to others. The early symptoms of scarlet fever include sore throat, headache, fever, nausea and vomiting. After 12 to 48 hours, a fine red rash develops which typically first appears on the chest and stomach, rapidly spreading to other parts of the body. The skin can feel a bit like sandpaper. The scarlet rash may be harder to spot on darker skin, although the 'sandpaper' feel should be present. Children often have flushed cheeks and may have a bright red 'strawberry' tongue.

If you think you, or your child, might have scarlet fever:

- contact your GP or NHS 111 as soon as possible
- make sure that you or your child take(s) the full course of any antibiotics prescribed. Although you or your child will feel better soon after starting the course of antibiotics, you must complete the course to ensure that you do not carry the bacteria in your throat after you have recovered
- stay at home, away from nursery, school or work for at least 24 hours after starting the antibiotic treatment, to avoid spreading the infection

You can help stop the spread of infection through frequent hand washing and by not sharing eating utensils, clothes, bedding and towels. All contaminated tissues should be disposed of immediately.

Stop the spread

During periods of high incidence of scarlet fever, there may also be an increase in outbreaks in schools, nurseries and other childcare settings. Children and adults with suspected scarlet fever should stay off nursery / school / work until **24 hours** after the start of appropriate antibiotic treatment. Good hygiene practice such as hand washing remains the most important step in preventing and controlling spread of infection.

Yours sincerely,

Louisa Hedley
Executive Assistant